

## Mercury Poisoning in a Free-Living Northern River Otter (*Lontra canadensis*)

Jonathan M. Sleeman,<sup>1,5,6</sup> Daniel A. Cristol,<sup>2</sup> Ariel E. White,<sup>2</sup> David C. Evers,<sup>3</sup> R. W. Gerhold,<sup>4</sup> and Michael K. Keel<sup>4</sup> <sup>1</sup> Virginia Department of Game and Inland Fisheries, Richmond, Virginia 23230, USA; <sup>2</sup> Department of Biology, The College of William & Mary, Williamsburg, Virginia 23187, USA; <sup>3</sup> BioDiversity Research Institute, Gorham, Maine 04038, USA; <sup>4</sup> Southeastern Cooperative Wildlife Disease Study, College of Veterinary Medicine, The University of Georgia, Athens, Georgia 30602, USA; <sup>5</sup> Current address: USGS National Wildlife Health Center, 6006 Schroeder Road, Madison, Wisconsin 53711, USA; <sup>6</sup> Corresponding author (email: jsleeman@usgs.gov)

**ABSTRACT:** A moribund 5-year-old female northern river otter (*Lontra canadensis*) was found on the bank of a river known to be extensively contaminated with mercury. It exhibited severe ataxia and scleral injection, made no attempt to flee, and died shortly thereafter of drowning. Tissue mercury levels were among the highest ever reported for a free-living terrestrial mammal: kidney, 353 µg/g; liver, 221 µg/g; muscle, 121 µg/g; brain (three replicates from cerebellum), 142, 151, 151 µg/g (all dry weights); and fur, 183 µg/g (fresh weight). Histopathologic findings including severe, diffuse, chronic glomerulosclerosis and moderate interstitial fibrosis were the presumptive cause of clinical signs and death. This is one of a few reports to document the death of a free-living mammal from presumed mercury poisoning.

**Key words:** *Lontra canadensis*, mercury poisoning, northern river otter, Virginia.

The accumulation of mercury in northern river otters (*Lontra canadensis*) is well documented (Mierle et al., 2000; Ben-David et al., 2001; Yates et al., 2005), and has been suggested as a cause of population declines (e.g., Hyvarinen et al., 2003) or reduced survivorship (Mierle et al., 2000). Mercury intoxication can cause behavioral aberrations in both American mink (*Neovison vison*) and otters, including circling and attempts to burrow into the ground (Wobeser et al., 1976; Wren, 1985). Experimental studies have been used to determine the possible level at which mercury results in toxic effects in these species by dosing animals with methylmercury (e.g., O'Connor and Nielsen, 1980). However, documented cases of fatal mercury accumulation in free-ranging wildlife are rare. The use of otters and mink as biomonitors for ecosystems is

increasing (Basu et al., 2007; Wolfe et al., 2007). Background levels of mercury in otters in ecosystems contaminated primarily through atmospheric deposition are under 9.0 µg/g (wet weight [ww]) in the liver (Yates et al., 2005) and lower in the kidney (Kucera, 1983). Levels of mercury in ecosystems contaminated with point sources can result in mortalities, as reported for a dead otter from Ontario with hepatic mercury levels of 96.0 µg/g, ww (Wren, 1985). The fortuitous discovery of a moribund animal in the South River, Shenandoah Valley, Virginia, a site where invertebrates, fish, and birds are known to be extensively contaminated with industrial mercury (Cristol et al., 2008), provided a rare opportunity to associate specific tissue levels with mortality in a wild otter. Analyses revealed the highest levels of mercury reported in otters.

In April 2006, a moribund northern river otter was observed on the western bank of the South River (38°12.531'W, 78°50.477'N), Augusta County, Virginia. Visual examination of the animal from close range revealed tremors, bilateral scleral injection, and marked ataxia. The animal made no attempt to flee when approached and eventually returned to the water and drowned within minutes. The carcass was collected and refrigerated for 18 hr until a gross necropsy could be performed. Samples of ovary (with follicle), liver, lung, kidney, and heart were placed in 10% buffered formalin. These tissues as well as muscle (biceps femoralis), bile, blood, and fur from the hind leg were collected and frozen at -30 C. The

rest of the carcass, including the head, was also stored in a secure freezer at  $-30$  C. The head was taken to the Virginia Department of Agriculture and Consumer Services Animal Health Laboratory, Harrisonburg, Rockingham County, Virginia, where the brain was removed for rabies testing at the Division of Consolidated Laboratory Services, Richmond, Virginia, and three samples of cerebellum were retained frozen for mercury analysis. A tooth was removed for cementum age analysis (Matson's Laboratory LLC, 8140 Flagler Road, Milltown, Montana, USA).

Tissues in 10% buffered formalin were submitted to Southeastern Cooperative Wildlife Disease Study, College of Veterinary Medicine, University of Georgia, Athens, Georgia, USA for histopathologic evaluation. Tissue sections were embedded in paraffin, sectioned at  $3\ \mu\text{m}$ , and stained with hematoxylin and eosin and Ziehl-Neelsen acid-fast for light microscopy.

Analysis for total mercury was completed with cold vapor atomic absorbance spectroscopy with the use of a direct mercury analyzer (DMA-80 Milestone, Inc., Shelton, Connecticut, USA) at the Trace Element Research Laboratory (Texas A&M University, College Station, Texas, USA). The factory-calculated instrument detection limit for the direct mercury analyzers used was  $0.005\ \text{ng}$  and was calculated to be between  $0.013$  and  $0.026$  ( $n=9$ ) during this analysis. Mean percent recovery of standard reference materials (DORM-2, DOLT-3; National Research Council Canada, Ottawa, Ontario, Canada) was  $100.75 \pm 4.79\%$  ( $n=4$ ) during the running of the samples reported while spike recovery was  $100.70 \pm 1.37\%$  ( $n=13$ ) for the week of analysis.

No gross lesions were observed on necropsy, except for lack of body fat. Histopathologically, the mesangium of the glomeruli was thickened by fibrous connective tissue. Fibrous connective tissue also expanded the cortical interstitium and to a lesser extent the medullary intersti-




FIGURE 1. Histopathologic section of a kidney of a river otter (*Lontra canadensis*) with presumed mercury poisoning. Note the thickening of the mesangium of the glomeruli by fibrous connective tissue with expansion into the cortical interstitium, and to a lesser extent the medullary interstitium. H&E stain. Bar= $20\ \mu\text{m}$ .

tium (Fig. 1). The liver contained a few to moderate numbers of lymphocytes and plasma cells and rare neutrophils that surrounded periportal areas. Similar infiltrates surrounded bronchioles in the lung. Microscopic lesions were not apparent in the heart or reproductive tract. The brain was immunonegative for rabies by fluorescent antibody and the lung and liver were immunonegative for canine distemper virus (CDV). The otter was determined to be 5 yr of age and the cementum annuli were clear and distinct.

Mercury concentration of tissues was (dry weight, except for fur): fur,  $183\ \mu\text{g/g}$  (fresh weight); kidney,  $353\ \mu\text{g/g}$  (75.1% moisture); liver,  $221\ \mu\text{g/g}$  (74.1% moisture); muscle,  $121\ \mu\text{g/g}$  (74.6% moisture), brain (3 replicates from cerebellum),  $142$  (78% moisture),  $151$  (78.2% moisture),  $151\ \mu\text{g/g}$  (74.1% moisture).

The clinical, pathologic, and toxicologic findings are consistent with the few previously published case reports of mercury poisoning in otters (Heinz, 1996). The lesions seen in the kidneys were severe and chronic and most likely led to the clinical signs in this otter; however, as brain tissue was not available for histopathology, other causes of the central

TABLE 1. Mercury levels reported in tissues from various mammal species.

Species	Location	Level ( $\mu\text{g/g}$ ) (dry weight)	Tissue	Reference
Otter	Clay Lake, Ontario, Canada	96.0	Liver	Wren (1985)
		58.0	Kidney	
		36.0	Muscle	
		30.0	Brain	
Otter	Experimentally killed	33.4	Liver	O'Connor and Nielsen (1980)
		39.2	Kidney	
		15.7	Muscle	
		18.9	Brain	
Mink	South Saskatchewan River, Saskatoon, Canada	58.2	Liver	Wobeser and Swift (1976)
		31.9	Kidney	
		15.2	Muscle	
		13.4	Brain	
		34.9	Fur	
Feral cats	Minamata Bay, Japan	37–145	Liver	Takeuchi et al. (1977)
Striped dolphin	French Mediterranean Coast	68–2,272	Liver	Augier et al. (1993)
		14–34	Kidney	
		7–155	Muscle	
		4–81	Brain	
Florida panther	Everglades, Florida, USA	330	Liver	Dunbar (1994) <sup>a</sup>

<sup>a</sup> Originally reported as wet weight. For the purposes of comparison we assumed that wet weight value was equal to dry weight value/3 (Puls, 1994).

nervous system signs such as encephalitis due to toxoplasmosis cannot be ruled out (Fernandez-Moran, 2003). However, the mercury concentration in the brain of this animal was markedly higher than previously reported levels in naturally and experimentally exposed otters (142–151  $\mu\text{g/g}$  versus 30.0 and 18.9  $\mu\text{g/g}$ , respectively) (O'Connor and Nielsen, 1980; Wren, 1985), suggesting mercury poisoning as the proximate cause of death.

Although the most consistent effects of methylmercury are on the central nervous system, renal necrosis and fibrosis have been reported in several experimentally dosed species, including guinea pigs, mink, and harp seals (*Phoca groenlandicus*), as well as in a naturally exposed horse (Heinz, 1996). The etiology of the renal pathologic changes is thought to be autoimmune glomerulopathy due to chronic inorganic mercury filtration (Eto et al., 1997). The remarkable finding in this case was the extremely high body burdens of mercury. With the possible exception of a Florida panther (*Felis*

*concolor coryi*), these levels are higher than previously reported values from other nonmarine mammals (Table 1), and set an upper boundary for documented lethal levels of mercury in otters. To the authors' knowledge this case report is the first to describe histopathologic changes associated with markedly elevated tissue mercury levels in a naturally exposed river otter.

The value of the river otter as an established indicator species for sublethal effects of mercury is recognized by national mercury monitoring programs (Wolfe et al., 2007); however, it has not been well documented that otters or other animals might suffer mortality from direct exposure to mercury pollution. Although sublethal and reproductive-effect thresholds are increasingly being identified in free-living piscivorous birds and mammals (Scheuhammer et al., 2007; Burgess and Meyer, 2008; Evers et al., 2008; Scheuhammer et al., 2008; Basu and Head, 2010), evidence of mortality related to mercury contamination in ecosystems, such as this account, is rare.

We thank V. Weiss for veterinary assistance. The otter was collected using Virginia Department of Game and Inland Fisheries Salvage/Collecting Permit 026945.

#### LITERATURE CITED

- AUGIER, H., W. K. PARK, AND C. RONNEAU. 1993. Mercury contamination of the striped dolphin (*Stenella coeruleoalba meyen*) from the French Mediterranean coasts. *Marine Pollution Bulletin* 26: 306–311.
- BASU, N., A. M. SCHEUHAMMER, S. J. BURSIA, J. ELLIOTT, K. ROUVINEN-WATT, AND H. M. CHAN. 2007. Mink as a sentinel species in environmental health. *Environmental Research* 103: 130–144.
- , AND J. HEAD. 2010. Mammalian wildlife as complementary models in environmental neurotoxicology. *Neurotoxicology and Teratology* 32: 114–119.
- BEN-DAVID, M., L. K. DUFFY, G. M. BLUNDELL, AND R. T. BOWYER. 2001. Natural exposure of coastal river otters to mercury: Relation to age, diet, and survival. *Environmental Toxicology and Chemistry* 20: 1986–1992.
- BURGESS, N. M., AND M. W. MEYER. 2008. Methylmercury exposure associated with reduced productivity in common loons. *Ecotoxicology* 17: 83–91.
- CRISTOL, D. A., R. L. BRASSO, A. M. CONDON, R. E. FOVARGUE, S. L. FRIEDMAN, K. K. HALLINGER, A. P. MONROE, AND A. E. WHITE. 2008. The movement of aquatic mercury through terrestrial food webs. *Science* 320: 335.
- DUNBAR, M. R. 1994. Florida panther biomedical investigation. Florida Game and Fresh Water Fish Commission, Gainesville, Florida, pp. 1–54.
- ETO, K., A. YASUTAKE, K. MIYAMOTO, H. TOKUNAGA, AND Y. OTSUKA. 1997. Chronic effects of methylmercury in rats. II. Pathological aspects. *Tohoku Journal of Experimental Medicine* 182: 197–205.
- EVERS, D. C., L. SAVOY, C. R. DESORBO, D. YATES, W. HANSON, K. M. TAYLOR, L. SIEGEL, J. H. COOLEY, M. BANK, A. MAJOR, K. MUNNEY, H. S. VOGEL, N. SCHOCH, M. POKRAS, W. GOODALE, AND J. FAIR. 2008. Adverse effects from environmental mercury loads on breeding common loons. *Ecotoxicology* 17: 69–81.
- FERNANDEZ-MORAN, J. 2003. Mustelidae. In *Zoo and wild animal medicine*, 5th Edition, M. E. Fowler and R. E. Miller (eds.). Saunders, St. Louis, Missouri, pp. 501–516.
- HEINZ, G. H. 1996. Mercury poisoning in wildlife. In *Noninfectious diseases of wildlife*, 2nd Edition, A. Fairbrother, L. N. Locke and G. L. Hoff (eds.). Iowa State University Press, Ames, Iowa, pp. 118–127.
- HYVARINEN, H., P. TYNI, AND P. NIEMINEN. 2003. Effects of moult, age, and sex on the accumulation of heavy metals in the otter (*Lutra lutra*) in Finland. *Bulletin of Environmental Contamination and Toxicology* 70: 278–284.
- KUCERA, E. 1983. Mink and otter as indicators of mercury in Manitoba waters. *Canadian Journal of Zoology* 61: 2250–2256.
- MIERLE, G., E. M. ADDISON, K. S. MACDONALD, AND D. G. JOACHIM. 2000. Mercury levels in tissues of otters from Ontario, Canada: Variation with age, sex, and location. *Environmental Toxicology and Chemistry* 19: 3044–3051.
- O'CONNOR, D. J., AND S. W. NIELSEN. 1980. Environmental survey of methylmercury levels in wild mink (*Mustela vison*) and otter (*Lutra canadensis*) from the northeastern United States and experimental pathology of methylmercurialism in the otter. In *Proceedings of the world furbearer conference*, Frostburg, Maryland, J. A. Chapman and D. Pursley (eds.), pp. 1728–1745.
- PULS, R. 1994. Mineral levels in animal health diagnostic data, 2nd Edition. Sherpa International, Clearbrook, British Columbia, Canada, 356 pp.
- SCHEUHAMMER, A. M., M. W. MEYER, M. B. SANDHEINRICH, AND M. W. MURRAY. 2007. Effects of environmental methylmercury on the health of wild birds, mammals, and fish. *Ambio* 36: 12–18.
- , N. BASU, N. BURGESS, J. E. ELLIOTT, G. D. CAMPBELL, M. WAYLAND, L. CHAMPOUX, AND J. RODRIGUE. 2008. Relationships among mercury, selenium, and neurochemical parameters in common loons (*Gavia immer*) and bald eagles (*Haliaeetus leucocephalus*). *Ecotoxicology* 17: 93–101.
- TAKEUCHI, T., F. M. D'ITRI, P. V. FISCHER, C. S. ANNETT, AND M. OKABE. 1977. The outbreak of Minamata disease (methyl mercury poisoning) in cats on Northwestern Ontario reserves. *Environmental Research* 13: 215–218.
- WOBESER, G., N. O. NIELSEN, AND B. SCHIEFER. 1976. Mercury and mink. II. Experimental methylmercury intoxication. *Canadian Journal of Comparative Medicine* 40: 34–45.
- , AND M. SWIFT. 1976. Mercury poisoning in a wild mink. *Journal of Wildlife Diseases* 12: 335–340.
- WOLFE, M. F., T. ATKESON, W. BOWERMAN, K. BURGER, D. C. EVERS, M. W. MURRAY, AND E. ZILLIOUX. 2007. Wildlife indicators. In *Ecosystem response to mercury contamination: Indicators of change*, R. Harris, D. P. Krabbenhoft, R. Mason, M. W. Murray, R. Reash and T. Saltman (eds.). SETAC, CRC Press, Webster, New York, pp. 123–189.
- WREN, C. D. 1985. Probable case of mercury

poisoning in wild otter, *Lutra canadensis*, in northwestern Ontario. Canadian Field-Naturalist 99: 112–114.

YATES, D. E., D. T. MAYACK, K. MUNNEY, D. C. EVERS, A. MAJOR, T. KAUR, AND R. J. TAYLOR. 2005. Mercury levels in mink (*Mustela vison*) and river otter (*Lontra canadensis*) from north-

eastern North America. Ecotoxicology 14: 263–274.

*Submitted for publication 16 November 2009.*  
*Accepted 10 March 2010.*